

Density of states in d -wave superconductors disordered by extended impurities

İnanç Adagideli,^{1,*} Daniel E. Sheehy,^{2,†} and Paul M. Goldbart^{3,‡}

¹*Instituut-Lorentz, Universiteit Leiden, Niels Bohrweg 2, Leiden, NL-2333 CA, The Netherlands*

²*Department of Physics & Astronomy, University of British Columbia, 6224 Agricultural Rd., Vancouver, BC, Canada V6T1Z1*

³*Department of Physics, University of Illinois at Urbana-Champaign, 1110 West Green Street, Urbana, Illinois 61801*

(Received 12 July 2002; published 31 October 2002)

The low-energy quasiparticle states of a disordered d -wave superconductor are investigated theoretically. A class of such states, formed via tunneling between the Andreev bound states that are localized around extended impurities (and result from scattering between pair-potential lobes that differ in sign), is identified. Its (divergent) contribution to the total density of states is determined by taking advantage of connections with certain one-dimensional random tight-binding models. The states under discussion should be distinguished from those associated with nodes in the pair potential.

DOI: 10.1103/PhysRevB.66.140512

PACS number(s): 74.25.Jb, 74.72.-h

I. INTRODUCTION

In recent years, considerable attention has been focused on the low-energy electron-hole quasiparticle spectral properties of the cuprate superconductors in the presence of impurity scattering. Much of impetus for this effort has its origin in the fact that many cuprate superconductors are randomly chemically doped insulators, and are therefore disordered. Moreover, as they are pair breakers for them, the role of impurities is especially important for d -wave superconductors. Of particular interest is the behavior of the single-particle density of states (DOS) $\rho(E)$ as the energy E tends to zero, i.e., its low-energy behavior.

In recent work on the DOS of disordered d -wave superconductors, Pépin and Lee¹ invoked a t -matrix approximation to infer that $\rho(E) \sim 1/E |\ln E^2|^2$ at low energies. More recently, Yashenkin *et al.*² and Altland³ argued that the divergence found in Ref. 1 is present only for the case of a vanishing chemical potential (i.e., for a half-filled band), and thus does not apply to a doped cuprate. (To be precise, unitarity of the impurity scattering is also required.) It was further argued in Refs. 2 and 3 that, instead of diverging, $\rho(E)$ should vanish at $E=0$ (unless certain very specific fine-tuning requirements are met). An important feature shared by Refs. 1–3 is the hypothesis that the disorder potential may be adequately modeled by a random collection of *pointlike* scatterers. However, for a *single* impurity in a d -wave superconductor, the low-energy DOS is qualitatively different for point-like⁴ and extended⁵ (i.e., impurities of a size much larger than the Fermi wavelength) impurities: the states that reside at zero energy for extended impurities reside at non-zero energies for pointlike impurities. (The underlying reason for this difference is that, for pointlike impurities, the quasiparticle scattering is essentially diffractive, whereas for the extended impurities it is essentially semiclassical.) This observation raises the possibility that such differences will continue to manifest themselves in the many-impurity setting.

The purpose of the present paper is to identify a mechanism for producing low-energy quasiparticle states. This mechanism is based on impurity-scattering processes that connect states associated with differing signs of the d -wave

pair potential.⁶ In the case of a single impurity,⁵ this mechanism has already been shown to produce low-energy states that are localized near the impurity. These states can be associated with the classical trajectories scattering from the impurity, and have been observed via scanning-tunneling spectroscopy.^{7,8}

Here we build upon this single-impurity physics to identify a singular (and potentially dominant) contribution to the low-energy DOS. This contribution, which conventional techniques fail to capture, arises from tunneling along the classical trajectories that connect the individual impurities and, hence, connect the low-energy states localized near these impurities. The underlying physics was formulated some time ago in the context of tunnelling corrections to ground-state energies in models of supersymmetric quantum mechanics.^{9–11}

The picture we have in mind (see Fig. 1) of the processes that lead to low-energy states involves classical trajectories that each visit many extended impurities. As a quasiparticle progresses along such a trajectory, its momentum is repeatedly altered via scattering from the extended impurities, so that the effective pair potential [Fig. 1(b)] undergoes sign changes. Localized near each such sign change would be a zero-energy quasiparticle state; quasiparticle tunneling (through the pair potential) connects these states, raising their energies from zero, and thus forming a low-energy band that exhibits a Dyson-like singularity¹² at zero energy: $\rho(E) \sim 1/E |\ln E^2|^3$. This picture loses its precision for sequences of impurities between which the pair potential is predominantly small (i.e., for nodal directions) and, consequently, the states are not well localized near the impurities. However, the contribution on which we are focusing (i.e., the non-nodal contribution) is expected to be substantial, and quite likely dominant, in the low-energy limit.

Based on previous work on low-energy quasiparticle states,^{5,4} we identify three classes of processes that can effect the low-energy DOS of a disordered d -wave superconductor: (i) purely semiclassical scattering between states with differing signs of the pair-potential (i.e., scattering due to random extended scatterers in the d -wave superconductors), which we shall focus on in the present paper; (ii) purely diffractive scattering between states near the nodal directions of the




FIG. 1. (a) Sketch of a classical trajectory that encounters several impurities and, hence, several sign changes in Δ . The shaded regions on the trajectory denote the sections of the trajectory where approximate zero-energy states reside. (b) Momentum-space pair potential of a d -wave superconductor. (c) Schematic depiction of Δ and Δ^2 along the trajectory.

pair-potential (i.e. the scattering of nodal quasiparticles by pointlike impurities), which was considered in, e.g., Refs. 3, 2, and 13; and (iii) the mixing of the low-energy states that arise from processes (i) and (ii). Throughout the rest of this paper we shall ignore processes in class (iii). In consequence, the low-energy DOS of a disordered d -wave superconductor can be expressed as a sum of contributions arising from class (i) processes (viz. ρ_{ext}) and class (ii) processes (viz. ρ_{nodal}), the latter, as discussed in Refs. 2 and 3, being nondivergent and therefore subdominant. Thus, we shall focus on ρ_{ext} [and denote this by $\rho(E)$].

II. QUASIPARTICLE MOTION IN A d -WAVE SUPERCONDUCTOR WITH MANY EXTENDED IMPURITIES

Our focus will be on the DOS per unit area, i.e.,

$$\rho(E) \equiv \frac{1}{A} \sum_m \delta(E - E_m), \quad (1)$$

where A is the area of the sample and the energy eigenvalues $\{E_m\}$ follow from the Bogoliubov–de Gennes eigenproblem, viz.,

$$\begin{pmatrix} \hat{h} & \hat{\Delta} \\ \hat{\Delta} & -\hat{h} \end{pmatrix} \begin{pmatrix} u_n \\ v_n \end{pmatrix} = E_n \begin{pmatrix} u_n \\ v_n \end{pmatrix}. \quad (2)$$

Here $\hat{h} \equiv -\nabla^2 - k_F^2 + V(\mathbf{r})$, in which k_F^2 is the chemical potential [i.e., k_F ($\equiv 2\pi/\lambda_F$) is the Fermi wave vector], V is the single-particle impurity potential, and we have adopted units in which $\hbar^2/2m = 1$, m being the common (effective) mass of the electrons and holes. The operator $\hat{\Delta}$ is the pair-

potential (integral) operator; how it acts is specified by the nonlocal pair-potential kernel $\Delta(\mathbf{r}, \mathbf{r}')$ via $[\hat{\Delta}v_n](\mathbf{r}) = \int d^2r' \Delta(\mathbf{r}, \mathbf{r}')v_n(\mathbf{r}')$.

To define this model fully, we need an assumption about the form of $\Delta(\mathbf{r}, \mathbf{r}')$. It is convenient to exchange the coordinates \mathbf{r} and \mathbf{r}' for the relative and center-of-mass coordinates $\boldsymbol{\rho}$ and \mathbf{R} :

$$\bar{\Delta}(\boldsymbol{\rho}, \mathbf{R}) \equiv \Delta(\mathbf{r}, \mathbf{r}'), \quad \boldsymbol{\rho} \equiv \mathbf{r} - \mathbf{r}', \quad 2\mathbf{R} \equiv \mathbf{r} + \mathbf{r}'. \quad (3)$$

Then, via Fourier transformation with respect to $\boldsymbol{\rho}$, viz.,

$$\bar{\Delta}(\mathbf{k}, \mathbf{R}) \equiv \int d^2\rho e^{-i\mathbf{k}\cdot\boldsymbol{\rho}} \bar{\Delta}(\boldsymbol{\rho}, \mathbf{R}), \quad (4)$$

we obtain the pair potential $\bar{\Delta}(\mathbf{k}, \mathbf{R})$ at the center-of-mass position \mathbf{R} and relative momentum \mathbf{k} . As our aim is to describe the cuprate superconductors, we take $\bar{\Delta}(\mathbf{k}, \mathbf{R})$ to have d -wave symmetry, $\bar{\Delta}(\mathbf{k}, \mathbf{R}) \propto (k_x^2 - k_y^2)$, where k_x and k_y denote the Cartesian components of \mathbf{k} . However, we allow for the possibility of position-dependent amplitude variations of the d -wave pair potential due, say, to pair-breaking effects near the extended scatterers.

III. SEMICLASSICAL APPROACH TO THE BOGOLIUBOV–DE GENNES EIGENPROBLEM

We now invoke a semiclassical approximation under which $\rho(E)$ is expressed in terms of the solution of a family of one-dimensional eigenproblems, each associated with a classical scattering trajectory in the presence of the single-particle impurity potential $V(\mathbf{r})$. We restrict ourselves to a brief discussion of this approach; for details, see Refs. 5 and 14. The approximation amounts to our: (i) regarding the kinetic and potential energies as being comparable and being the largest energies in the problem, (ii) turning off the pair potential, (iii) semiclassically treating the quasiparticle motion in the presence of the kinetic and potential energies, and (iv) reinstating the pair potential. Via this approach, we reduce the two-dimensional Bogoliubov–de Gennes eigenproblem to a family of one-dimensional Andreev eigenproblems residing on trajectories, each trajectory being a particular classical scattering trajectory in the presence of the many-impurity potential. This scheme applies under the following conditions: (i) the amplitudes of $\bar{\Delta}$ and V should vary slowly, relative to the Fermi wavelength λ_F ; and (ii) the Fermi energy k_F^2 should be large compared with the energy scale of interest, viz. E , as well as with the typical pair-potential scale.

Let us now turn to the family of one-dimensional eigenproblems arising from this semiclassical scheme. Following Ref. 5, these trajectory-dependent eigenproblems take the form

$$\hat{H} \begin{pmatrix} \bar{u}_n \\ \bar{v}_n \end{pmatrix} = E_n \begin{pmatrix} \bar{u}_n \\ \bar{v}_n \end{pmatrix}, \quad (5a)$$

$$\hat{H} \equiv \begin{pmatrix} -2ik_F \partial_\sigma & \Delta_0(\sigma) \\ \Delta_0(\sigma) & 2ik_F \partial_\sigma \end{pmatrix}, \quad (5b)$$

$$\Delta_0(\sigma) \simeq \bar{\Delta}[k_F \partial_\sigma \mathbf{x}_c(\sigma), \mathbf{x}_c(\sigma)], \quad (5c)$$

i.e., an Andreev eigenproblem.¹⁵ Here, the parameter σ measures the position along a particular classical trajectory $\mathbf{x}_c(\sigma)$, the latter obeying Newton's equation in the many-impurity potential, viz.,

$$k_F^2 \partial_\sigma^2 \mathbf{x}_c(\sigma) = -\nabla V[\mathbf{x}_c(\sigma)]. \quad (6)$$

The DOS is then obtained by assembling the eigenvalue spectra $E_n(\mathbf{n}, b)$ of all the classical trajectories, the latter being labeled in terms of an asymptotic momentum direction \mathbf{n} and impact parameter b :

$$\rho(E) \simeq \frac{k_F}{A} \int \frac{d\mathbf{n}}{2\pi} \int db \rho(\mathbf{n}, b, E), \quad (7a)$$

$$\rho(\mathbf{n}, b, E) \equiv \sum_m \delta[E - E_m(\mathbf{n}, b)]. \quad (7b)$$

Thus, in order to obtain $\rho(E)$ one needs to find each classical trajectory, obtain the associated effective pair-potential [given by Eq. (5c)], solve the resulting one-dimensional eigenvalue equation, and, finally, integrate over all the classical trajectories using Eq. (7a). We note that if one interprets the weight of a particular classical trajectory as the probability of finding a pair-potential configuration corresponding to that particular trajectory, then we see that the calculation of $\rho(E)$ amounts to computing the average density of states of a random pair-potential model. Models of this sort were considered, e.g., in Refs. 16–19.

IV. EIGENVALUE PROBLEM FOR A SINGLE TRAJECTORY

We now examine the contribution $\rho(\mathbf{n}, b, E)$ to the DOS for the case of a generic trajectory (\mathbf{n}, b) . For convenience, we introduce the rescaled trajectory parameter $s \equiv \sigma/2k_F$; the Hamiltonian then becomes

$$\hat{H} = \begin{pmatrix} -i\partial_s & \Delta(s) \\ \Delta(s) & i\partial_s \end{pmatrix}, \quad (8)$$

where $\Delta(s) \equiv \Delta_0(2k_F s)$.

Our method for calculating the spectrum of \hat{H} in the many-impurity case is based on that for a single impurity.⁵ In the latter case, low-energy states arose from asymptotically sign-changing trajectories [i.e. those trajectories for which $\lim_{s \rightarrow \pm\infty} \Delta(s)$ differ in sign]. Finding the spectrum amounted to identifying such sign-changing trajectories. What about the case of *many* extended impurities? In this case, for a typical trajectory through the impurity potential $\Delta(s)$ undergoes repeated sign changes. On a particular trajectory let us label the positions of these zeroes of $\Delta(s)$ by $\{s_n\}$. Recall that we are concerned with the collection of impurity states that would lie at zero energy if the impurities were isolated. Owing to tunneling between them, these formerly

degenerate states yield a continuum of states, extending upward in energy from zero. Our task is to shed some light on this band formation. We proceed to set up a tight-binding model along the trajectory, in which we retain only the zero-energy impurity states $\{|n\rangle\}$ (i.e., the local ground states at each of the $\{s_n\}$) and allow only nearest-neighbor tunneling between them.^{20,21} To complete the model, we need the matrix elements of \hat{H} connecting these states, i.e.,

$$t_n \equiv \langle n | \hat{H} | n+1 \rangle. \quad (9)$$

Using the analytic expression for the zero energy wave functions,¹¹ $\langle s | n \rangle \propto \exp -\int_{s_n}^s ds' |\Delta(s')|$, a straightforward calculation produces

$$t_n \approx \frac{1}{\sqrt{\pi}} |\Delta'(s_n) \Delta'(s_{n+1})|^{1/4} \exp \left\{ - \int_{s_n}^{s_{n+1}} ds' |\Delta(s')| \right\}, \quad (10)$$

where $\Delta'(s) \equiv \partial_s \Delta(s)$. We are now in a position to write down a low-energy effective approximation to \hat{H} , viz.,

$$\hat{H} \approx \sum_n t_n (|n\rangle \langle n+1| + |n+1\rangle \langle n|), \quad (11)$$

i.e., for each classical trajectory, one arrives at a (topologically) one-dimensional hopping model that captures the physics of tunneling processes between the (formerly zero-energy) states localized near each zero of the pair-potential.

V. DENSITY OF STATES

In order to obtain the low-energy DOS, we must obtain the DOS of the effective Hamiltonian [Eq. (11)] for each trajectory, and then collect them together. We assume that the collection of trajectories forms an ensemble that is characterized by the condition that momentum directions before and after a collision are uncorrelated. Then, summing over such an ensemble of trajectories is equivalent (up to a constant of proportionality) to averaging the DOS of Hamiltonian (11) over uncorrelated values of t_n . To obtain the low-energy DOS of this effective model we appeal to results obtained by Eggarter and Riedinger,²² who, building on the work of Dyson¹² and Theodorou and Cohen,²³ studied random-hopping models of precisely this form. Specifically, in Ref. 22 it was found that, under the condition that the $\{t_n\}$ are uncorrelated²⁴ and identically distributed, the DOS as $E \rightarrow 0$ is given by

$$\rho(E) \approx N_s Z \frac{2\sigma^2}{E |\ln(E/\bar{t})|^2|^3}, \quad (12)$$

where N_s denotes the average number of sites along the trajectory, Z is the constant of proportionality arising from the Jacobian of the transformation from summing over trajectories to averaging over $\{t_n\}$, \bar{t} is the scale characterizing $\{t_n\}$, and the amplitude σ^2 is given by the variance of the logarithm of t , i.e.,

$$\sigma^2 \equiv \langle (\ln t^2 / \Delta_0^2)^2 \rangle - \langle \ln t^2 / \Delta_0^2 \rangle^2, \quad (13)$$

where $\langle \dots \rangle$ denotes a disorder average. The scales \bar{t} and σ^2 could be estimated with the help of Eq. (10) as

$$\bar{t}^2 \propto k_F \Delta_0 n_c^{1/2} \exp(-\Delta_0 / k_F n_c^{1/2}), \quad (14a)$$

$$\sigma^2 \propto \Delta_0 / k_F n_c^{1/2}, \quad (14b)$$

where n_c is the number of impurities per unit area. What remains is to determine the coefficient Z ; we now make an estimate of this quantity.

Our estimate for Z follows from considering the integration over b for a single impurity of size a . As the impurity potential is expected to decay rapidly away from the impurity, the only trajectories that interact appreciably with it are those that directly intersect it, i.e., $k_F \int db \rightarrow k_F a$. Thus, in the absence of the tunneling corrections between the zero-energy states, $\rho(E)$ is approximately given by $\rho_0(E) \approx k_F a n_c \delta(E)$.⁵ Although the inclusion of tunneling corrections changes the energy dependence of $\rho(E)$, here we assume that the energy-integrated density of states $\int_0^\epsilon \rho(E) dE$ is approximately conserved for some appropriately chosen cutoff ϵ . These considerations lead to the following approximate form for $\rho(E)$ valid for $E \ll \bar{t}$:

$$\rho(E) \propto \frac{\Delta_0 a n_c^{1/2}}{E |\ln(E/\bar{t})|^2 |^3}. \quad (15)$$

We remark that the divergence of the low-energy DOS is ultimately cut off due to physical processes not included in the present description. These include the dephasing scale \hbar/τ_ϕ and the diffractive scattering scale $\hbar/\tau_{d.s.}$; the cut-off will occur at the largest of these scales. Thus it is possible that at extremely low energies $\rho(E)$ eventually does vanish asymptotically, in agreement with the results of Refs. 2 and 3. However, the present results would still apply at intermediate energies. Finally, we stress that the calculation presented here does not place any special emphasis on the nodes of the d -wave order parameter; indeed, the singular contribution to the density of states of a d -wave superconductor reported here arises from states in generic (rather than nodal) regions on the Fermi surface.

ACKNOWLEDGMENTS

P.M.G. thanks Alexei Tselik for informative discussions and the University of Colorado at Boulder for its kind hospitality. This work was supported by the U.S. Department of Energy under Award No. DEFG02-ER9645439, through the Frederick Seitz Materials Research Laboratory, by the Dutch Science Foundation NWO/FOM (I.A.), and by NSERC (DES).

*Electronic address: adagidel@lorentz.leidenuniv.nl

†Electronic address: sheehy@physics.ubc.ca

‡Electronic address: goldbart@uiuc.edu

¹C. Pépin and P.A. Lee, Phys. Rev. B **63**, 054502 (2001).

²A.G. Yashenkin, W.A. Atkinson, I.V. Gornyi, P.J. Hirschfeld, and D.V. Khveshchenko, Phys. Rev. Lett. **86**, 5982 (2001).

³A. Altland, Phys. Rev. B **65**, 104525 (2002).

⁴A.V. Balatsky, M.I. Salkola, and A. Rosengren, Phys. Rev. B **51**, 15 547 (1995); M.I. Salkola, A.V. Balatsky, and D.J. Scalapino, Phys. Rev. Lett. **77**, 1841 (1996).

⁵I. Adagideli, P.M. Goldbart, A. Shnirman, and A. Yazdani, Phys. Rev. Lett. **83**, 5571 (1999).

⁶Of course, the mechanism for creating low-energy states invoked in the present paper does not specifically require d -wave superconductivity. All that is required is that the pair potential be real (or can be made so by a suitable gauge transformation) and that it possess regions of momentum orientation over which it has opposing signs.

⁷A. Yazdani, C.M. Howald, C.P. Lutz, A. Kapitulnik, and D.M. Eigler, Phys. Rev. Lett. **83**, 176 (1999).

⁸Similar states have also been observed in the case of native defects in E. Hudson, S.H. Pan, A.K. Gupta, K.-W. Ng, and J.C. Davis, Science **285**, 88 (1999).

⁹E. Witten, Nucl. Phys. B **188**, 513 (1981).

¹⁰P. Salomonson and J.W. Van Holten, Nucl. Phys. B **196**, 509 (1982).

¹¹See, e. g., G. Junker, *Supersymmetric Methods in Quantum and Statistical Physics* (Springer, Berlin, 1996).

¹²F.J. Dyson, Phys. Rev. **92**, 1331 (1953).

¹³C. Chamon and C. Mudry, Phys. Rev. B **63**, 100503 (2001).

¹⁴D.E. Sheehy, I. Adagideli, P.M. Goldbart, and A. Yazdani, Phys. Rev. B **64**, 224518 (2001).

¹⁵A.F. Andreev, Zh. Éksp. Teor. Fiz. **46**, 1823 1964 [Sov. Phys. JETP **19**, 1228 (1964)].

¹⁶D. Waxman and K.D. Ivanova-Moser, Ann. Phys. (N.Y.) **226**, 271 (1993).

¹⁷A. Comtet, J. Desbois, and C. Monthus, Ann. Phys. (N.Y.) **239**, 312 (1995).

¹⁸L. Balents and M.P.A. Fisher, Phys. Rev. B **56**, 12970 (1997).

¹⁹See, e. g., L. Bartosch, Ann. Phys. (Leipzig) **10**, 799 (2001).

²⁰Higher-order tunneling matrix elements are small, relative to the leading-order contributions, provided that $\int_{s_n}^{s_{n+1}} |\Delta(s)| ds \gg 1$ (Ref. 10), i. e., that the coherence length is small compared with the typical impurity spacing. In addition, we note that, as \hat{H} may be made purely off-diagonal by a suitable unitary transformation, the tunneling amplitude between next-nearest neighbors vanishes identically.

²¹The related task of computing tunneling corrections to the ground state of a SUSY Hamiltonian having multiple degenerate minima using instanton methods was considered in Ref. 10.

²²T.P. Eggarter and R. Riedinger, Phys. Rev. B **18**, 569 (1978).

²³G. Theodorou and M.H. Cohen, Phys. Rev. B **13**, 4597 (1976).

²⁴We suspect that the result for the DOS presented here will continue to hold not solely in the setting of uncorrelated hopping matrix elements $\{t_n\}$ but also for correlations of finite range.